

Bat Rabies in Georgia, 1956-65

JOHN H. RICHARDSON, D.V.M., M.P.H., RALPH L. RAMSEY, A.B., and L. E. STARR, D.V.M., Ph.D.

THE first laboratory-confirmed infection of rabies in an insectivorous bat was reported by Pawan (1) in a white bat, *Dictidurus alba*, in 1936. This white bat was among a group of insect-eating, fruit-eating, and blood-lapping species examined during the investigation of an outbreak of rabies in cattle and humans in Trinidad in 1932.

Rabies infection in an insectivorous bat from North America was first reported by Venters and associates (2) in a yellow bat, *Dasypterus floridanus*, killed near Tampa, Fla., in 1953. In the 13 years following (1953-65) rabies was diagnosed in one or more species of insectivorous bats from 47 States (personal communication, Dr. R. K. Sikes, Communicable Disease Center, 1966) and from two Provinces of Canada (3, 4).

Rabies infection in an insectivorous bat from Georgia was first reported in a Seminole bat, *Lasiurus seminobus*, found dead on a street in Thomas County, by personnel of the Public Health Service's Newton Field Station in 1956 (personal communication, Dr. Douglas Hawkins, Newton Field Station, 1956).

In the 10 years following (1956-65) rabies was diagnosed in six species of bats and from 18 counties in the State.

This paper reviews the 36 cases of laboratory-confirmed rabies in 531 insectivorous bats examined in Georgia during the 1956-65 period.

Materials and Methods

The 531 bats examined came from two primary sources. Residents of Georgia submitted 313 bats through routine diagnostic facilities. All of the 36 bats eventually diagnosed as rabid

were obtained this way. Two of these were initially taken to the Tennessee Department of Public Health Regional Laboratory in Chattanooga where the diagnosis was made. The carcasses were then forwarded to Atlanta for species identification and salivary gland examination (personal communication, L. V. Smith, Chattanooga Regional Laboratory, Tennessee Department of Public Health, 1965).

Field personnel collected the remaining 218 bats examined. Collections, confined almost exclusively to colonial species, were often prompted by complaints of bats inhabiting buildings, human exposures to bats, and discovery of infected bats in a county in which rabies had not previously been reported in these species.

Laboratory procedures. The initial rabies diagnostic procedure employed was microscopic examination of whole brain cross section impressions stained by Sellers' method. Negri-positive brain tissues were often injected intracerebrally into mice for additional diagnostic confirmation. Negri-negative brain tissues from bats that had bitten humans or had clinical signs suggestive of rabies were also selectively subjected to mouse inoculation tests.

When brain tissues of bats collected by field personnel between 1956 and 1960 proved to be Negri-negative, the tissues were often pooled (3-5 brains in each pool) for mouse inoculation tests. Since 1961 the fluorescent rabies

Dr. Richardson is public health veterinarian, Mr. Ramsey is public health ecologist, and Dr. Starr, now retired, was public health veterinarian, Georgia Department of Public Health, Atlanta.

antibody test has been used routinely in conjunction with microscopic examination and mouse inoculation of individual bat brain tissues in the diagnosis of rabies.

Intracerebral inoculation of mice was also used to demonstrate the presence of rabies virus in the salivary glands of selected specimens. No attempt was made to quantitate rabies virus in the brains or salivary glands of infected bats or to examine brown fat and other tissues for virus.

Species identification. Species identification of individual bats was based on gross physical characteristics including color, length, and distribution of pelage; relationship of tail and interfemoral membrane; dentition; and geographic distribution (5-8).

Bats whose size and dentition were within specified adult limits were designated as mature. Other specimens were designated as immature.

Results

A wide range of behavior was reported for the 36 rabid bats. Nineteen were found alive on the ground, apparently incapable of flight. These grounded bats were often aggressive, or perhaps defensive, when approached or manipulated. Six of the rabid bats were found dead on the ground; eight were either captured alive or brought dead into the yard by pet dogs or cats, and two made aggressive flying attacks on humans although no actual bites or contacts occurred. Two bats, a male and a female, were observed fighting in mid-air, captured after fall-

Table 1. Species distribution of rabies infection in 531 bats examined in Georgia, 1956-65

Species	Number examined	Number rabid	Percent rabid
Colonial bats.....	303	4	1.3
<i>Nycticeius humeralis</i> (twilight bat).....	172	2	1.2
<i>Tadarida cynocephala</i> (eastern free tailed bat).....	81	1	1.3
<i>Eptesicus fuscus</i> (big brown bat).....	32	0	0
<i>Pipistrellus subflavus</i> (southeastern pipistrelle bat).....	7	0	0
<i>Myotis austroriparius</i> (southeastern Myotis).....	6	1	16.6
<i>Myotis grisescens</i> (gray Myotis).....	3	0	0
<i>Myotis keenii</i> (Keen's Myotis).....	1	0	0
<i>Myotis</i> species.....	1	0	0
Noncolonial bats.....	228	32	14
<i>Lasiurus borealis</i> (red bat).....	173	26	15
<i>Lasiurus seminolus</i> (Seminole bat).....	27	3	11
<i>Lasiurus cinereus</i> (hoary bat).....	15	3	20
<i>Lasionycterus noctivagans</i> (silver haired bat).....	11	0	0
<i>Corynorhinus macrotis</i> (eastern big eared bat).....	2	0	0
Total.....	531	36	6.7

ing to the ground, and submitted for laboratory examination. Only the female was rabid.

The incidence of rabies, by species, in the 531 bats examined is given in table 1. The five colonial species accounted for 57 percent of the total examinations. However, the infection rate of the noncolonial bats was more than 10 times higher than that of the colonial bats.

Table 2. Rabies infection in 36 bats in Georgia by species, sex, and estimated stage of maturity, 1956-65

Species	Male		Female		Total
	Mature	Immature	Mature	Immature	
<i>Lasiurus borealis</i>	10	3	12	1	26
<i>Lasiurus cinereus</i>	1		2		3
<i>Lasiurus seminolus</i>			2		1 ¹ 3
<i>Nycticeius humeralis</i>	1		1		2
<i>Tadarida cynocephala</i>	1				1
<i>Myotis austroriparius</i>	1				1
Total.....	14	3	17	1	36


¹ Includes 1 specimen with no data concerning sex or maturity.

The species, sex, and estimated stage of maturity of each of the 36 rabid bats is given in table 2. Noncolonial bats of the genus *Lasiurus* accounted for 32 (89 percent) of the 36 confirmed infections. Within the lasiurine group the red bat, *Lasiurus borealis*, was found to be the species most frequently infected with rabies. Twenty-six (64 percent) of the rabid bats were red bats. Rabies infection was

equally distributed among male and female bats. Thirty-one of the 35 bats for which maturity data were available were mature specimens.

Golley (7) lists 9 genera and 15 species under the order *Chiroptera* reported to occur naturally in Georgia. Representatives of 12 of these 15 species were examined for rabies during the 10-year study period. Rabies was diagnosed

Figure 1. Geographic distribution of 36 cases of bat rabies in Georgia, 1956-65



in one or more individuals of 6 of these 12 species examined.

The geographic distribution of rabid bats is shown by species and county in figure 1. Approximately half of the cases were reported from the Metropolitan Atlanta area which consists in part of Clayton, Cobb, De Kalb, and Fulton Counties. The remaining cases were sporadically distributed in the Coastal Plain, Piedmont, and Ridge and Valley Provinces.

The monthly distribution of the 36 cases is shown in figure 2. On a calendar year basis the earliest reported case of bat rabies was on March 27 and the latest on November 5. More than 60 percent of the cases were reported during the 4-month period of July to October. A secondary peak of reported cases occurred in the spring.

The annual incidence of bat rabies showed considerable variation and was apparently influenced by both the species and number of bats examined in a given year. Reported cases ranged from a low of 1 rabid bat in 10 examined in 1959 to a high of 15 rabid bats in 159 examined in 1965.


Table 3 summarizes the results obtained in the laboratory examination of brain tissues and salivary glands of rabid bats by the applicable diagnostic procedure. Negri bodies were observed in 33 (94 percent) of the 35 brain tissues examined by Sellers' method. During the same 10-year period, Negri bodies were demonstrated in brain tissues of 323 (89 percent) of the 362 rabid dogs examined by Sellers' method.

Rabies virus was demonstrated, by mouse

Table 3. Results of laboratory examinations of tissues from 36 rabid bats in Georgia, 1956-65

Diagnostic procedure	Tissue examined	Number examined	Number positive for rabies
Microscopic examination.	Brain-----	35	33
Fluorescent rabies antibody.	-----do-----	25	25
Mouse inoculation-----	-----do-----	23	23
Do-----	Salivary glands.	21	12

Figure 2. Monthly distribution of 36 cases of bat rabies in Georgia, 1956-65



inoculation tests, in the salivary glands of 12 (57 percent) of the 21 rabid bats examined. Rabies antigen was demonstrated in all of the 25 rabid bats examined by the fluorescent rabies antibody procedure since 1961.

Discussion

Although the number of specimens described in this 10-year summary is limited, certain trends are apparent. Bat rabies in Georgia seems to be primarily associated with noncolonial adult lasiurine bats during the milder seasons of the year. This period of highest recognized incidence coincides with a period of relatively vigorous activity for most bats. In the early summer the rapidly developing young are weaned and become self-sufficient fliers and feeders. Late summer is associated with breeding activities, premigration wanderings, and migration in many species.

Reporting of bat rabies seems to be strongly influenced by the availability of interested observers. The large number of cases reported from urban areas suggests that the density of the human population may increase the opportunity of seeing a bat acting unusually. Announcement of cases in public media also increases general awareness and interest in bat rabies and stimulates the submission of specimens to the laboratory.

Rabies is thought to be transmitted in nature primarily by the bite of an infected animal.

The relatively high population density of colonial bats as compared to noncolonial bats should increase the opportunity for transmission of rabies by bite and nonbite (9) routes. The findings of this review and of previous investigations in Florida (10) do not corroborate this assumption. The rabies infection rate in noncolonial bats was found to be more than 10 times higher than in colonial bats in Georgia and more than 7 times higher in Florida.

Bat rabies in Georgia has occurred predominantly in counties that were free of reported rabies in terrestrial mammals. Only 7 of the 36 cases occurred in counties with rabies diagnosed in other animals during the same calendar year. No epidemiologic evidence exists to date which suggests the transmission of rabies by bats to other animals in Georgia.

The historical references cited by Pawan (1) indicate that vampire bats were associated with disease and death in South and Central America at the time of the Spanish colonization in the 18th century. The recently recognized and widespread distribution of rabid insectivorous bats in the United States is perhaps suggestive of a new ecological facet of the disease.

REFERENCES

- (1) Pawan, J. L.: The transmission of paralytic rabies in Trinidad by the vampire bat. *Ann Trop Med Parasit* 30: 101-131 (1936).
- (2) Venters, H. D., Hoffert, W. R., Scatterday, J. E., and Hardy, A. V.: Rabies in bats in Florida. *Amer J Public Health* 44: 182-185 (1954).
- (3) Avery, R. J., and Tailyour, J. M.: The isolation of the rabies virus from insectivorous bats in British Columbia. *Canad J Comp Med* 24: 143-145 (1960).
- (4) Beauregard, M., and Stewart, R. C.: Bat rabies in Ontario. *Canad J Comp Med* 28: 43-45 (1964).
- (5) Miller, G. S., and Kellogg, R.: List of North American recent mammals. U.S. National Museum Bulletin No. 205. U.S. Government Printing Office, Washington, D.C., 1955.
- (6) Hamilton, W. J.: The mammals of eastern United States. Comstock Publishing Co., Ithaca, N.Y., 1943.
- (7) Golley, F. B.: Mammals of Georgia. University of Georgia Press, Athens, 1962.
- (8) Hall, E. R., and Kelson, K. R.: The mammals of North America. Ronald Press Co., New York, 1959, vol. 1.
- (9) Constantine, D.: Rabies transmission by non-bite route. *Public Health Rep* 77: 287-289 (1962).
- (10) Schneider, N. J., et al.: Rabies in bats in Florida. *Amer J Public Health* 47: 983-989 (1957).

Experimental Vaccine Against Pneumonia

An experimental vaccine against primary atypical pneumonia that has given significant protection against experimentally produced illness has been tested by Dr. Charles B. Smith, Dr. William Friedewald, Dr. Robert Alford, and Dr. Robert M. Chanock of the National Institute of Allergy and Infectious Diseases (NIAID), Public Health Service. Results of the early trials were reported at the New York Academy of Science Conference on Mycoplasma, May 10-13, 1966.

The vaccine, which consists of a formalin-inactivated suspension of *Mycoplasma pneumoniae* organisms, was given to 19 volunteers who lacked prior antibody to the organism. Ten developed antibody and when these 10 were later experimentally infected with *M. pneumoniae*, only 1 became ill. Illness occurred in 10 of 13 control subjects who had not received the vaccine.

The killed vaccine, prepared by Chas. Pfizer & Co., Inc., is currently being tested on a larger scale in military populations under the sponsorship of NIAID's collaborative vaccine development program. The research group is also doing studies directed toward development of a live vaccine. *M. pneumoniae* infections are one of the greatest health problems in the armed services and are the cause of up to 50 percent of pneumonia cases in the 20-30 year age group in the general population.

PUBLICATION ANNOUNCEMENTS

Address inquiries to publisher or sponsoring agency.

Culture and Alcohol Use. A bibliography of anthropological studies. By Robert E. Popham and Carole D. Yawney. 1966; 47 pages. Alcoholism and Drug Addiction Research Foundation of Ontario, 24 Harbord St., Toronto 5, Canada.

Group Practice and Prepayment of Medical Care. By William A. MacColl, M.D. 1966; 257 pages; \$5. Group Health Association of America, 1321 14th St., NW., Washington, D.C. 20005.

The Potomac, People, Pollution. A report to the public. 1966; 30 pages. District of Columbia Department of Public Health, Health Education and Information Division, Washington, D.C. 20001.

Hospital Utilization Research Project, 1961-1966. Summary report supported by NIH Grant No. HM-00194. 1966; 64 pages. California Department of Public Health, Bureau of Hospitals, 2151 Berkeley Way, Berkeley, Calif. 94704.

Occupational Disease in California, 1964. By Diane Budde. 1966; 22 pages. California Department of Public Health, Bureau of Health Education, 2151 Berkeley Way, Berkeley, Calif. 94704.

Occupational Disease in California Attributed to Pesticides and Other Agricultural Chemicals, 1964. By Goldy D. Kleinman. 1966; 28 pages. California Department of Public Health, Bureau of Health Education, 2151 Berkeley Way, Berkeley, Calif. 94704.

Wages and Hours in the Retail Drug Industry, California—June 1965. 1966; 39 pages. State Department of Industrial Relations, Division of Labor Statistics and Research, 455 Golden Gate Ave., San Francisco, Calif.

Hospital Study Districts for Metropolitan Chicago. A geographic analysis and methodology. Techni-

cal Report No. 2. By Pierre de Visé. 1966; 75 pages. Hospital Planning Council for Metropolitan Chicago, 79 West Monroe St., Chicago, Ill. 60603.

Pattern for Change in Kentucky Mental Health Programs and Services. 1966; 233 pages. Kentucky Mental Health Planning Commission, Kentucky Department of Mental Health, P.O. Box 678, Frankfort, Ky.

Good Health. Personal and community. By Benjamin F. Miller, M.D., and John J. Burt, Ed.D. 2d edition. June 1966; 508 pages; \$7. W. B. Saunders Co., West Washington Square, Philadelphia, Pa. 19105.

Feasible Planning for Social Change. By Robert Morris and Robert Binstock. May 1966; 169 pages; \$5. Columbia University Press, 2660 Broadway, New York, N.Y. 10027.

Pediatric Nursing. By Audrey J. Kalafatic, R.N. June 1966; 432 pages; \$7. G. P. Putnam's Sons, 200 Madison Ave., New York, N.Y.

College Education. Key to a professional career in nursing. 1966; 15 pages; 25 cents. National League for Nursing, 10 Columbus Circle, New York, N.Y. 10019.

Master Education. Route to opportunities in modern nursing. 1966; 15 pages; 25 cents. National League for Nursing, 10 Columbus Circle, New York, N.Y. 10019.

Demand for Rehabilitation in a Labor Union Population. Action program. Pt. 2. By Hyman J. Weiner, D.S.W., Shelley H. Akabas, M.B.A., Morris Brand, M.D., and Bruce Grynbaum, M.D. 1966; 100 pages. Sidney Hillman Health Center, 16 East 16th St., New York, N.Y. 10003.

Death and Dying: Attitudes of patient and doctor. GAP Symposium No. 11. 1966; 80 pages; \$1.50. Group for the Advancement of Psychiatry, 104 East 25th St., New York 10010.

Further Studies of Automobile Glazing as an Injury Factor in Accidents. Automotive Crash Injury Research Cal No. VJ-1825-R3. March 1966; 77 pages. Cornell Aeronautical Laboratory, Inc. of Cornell University, Buffalo, N.Y. 14221.

A Method of Hospital Utilization Review. By Sidney Shindell, M.D., LL.B., and Morris London, M.P.H. 1966; 64 pages; \$1. University of Pittsburgh Press, Pittsburgh, Pa. 15213.

Successful Culturing of Glover's Cancer Organism and Development of Metastasizing Tumors in Animals Produced by Cultures from Human Malignancy. By George A. Clark, M.D., 1966; 37 pages. Cancer Research Foundation, 507 Liberty Ave., Pittsburgh, Pa.

Rehabilitative Aspects of Nursing. A programed instruction series. 1966; 51 pages; \$1.75. National League for Nursing, Research and Studies Service, 10 Columbia Circle New York 10019.

Fatal Home Accidents, Wisconsin, 1964. 1966; 11 pages. State Board of Health, P.O. Box 309, Madison, Wis. 53702.

The Lone Driver Involved in Injury-Producing Accidents. Automotive Crash Injury Research Cal No. VJ-1823-R19. By William E. Schotz. 1966; 49 pages. Cornell Aeronautical Laboratory, Inc. of Cornell University, P.O. Box 235, Buffalo, N.Y. 14221.

World Health Organization

WHO publications may be obtained from the Columbia University Press, International Documents Service, 2960 Broadway, New York, N.Y. 10027.

Sampling Methods in Morbidity Surveys and Public Health Investigations. WHO Expert Committee on Health Statistics. WHO Technical Report Series No. 336. 10th report. 1966; 29 pages; 60 cents; Geneva.